

The authors are grateful to the authorities in Spain who permitted entomological studies in their areas, to Dr H. Hacker (Staffelstein, Germany) for much helpful advice on the continental distribution of *L. nickerlii*, to Messrs J. Platts and J. Reid for their comments on aspects of the distribution and biology of the new subspecies, and to the Trustees of the Natural History Museum (London) for permission to study the collections there, and especially to Messrs D.J. Carter and M.R. Honey for their willing assistance.

#### References

- Culot, J., 1909-1913. *Les Noctuelles et Géomètres d'Europe*, Première Partie, Noctuelles 1: 220pp. Geneva.
- Draudt in Seitz, A., 1931-1938. Die Gross-Schmetterlinge der Erde, 3. *Supplement*, vi + 333pp. Stuttgart.
- Freyer, C.F., 1845. *Neuere Beitr. Schmett.* 5: 140; pl. 466, fig. 4.
- Ganev, J., 1982. Systematic and Synonymic List of Bulgarian Noctuidae (Lepidoptera). *Phegea* 10(3): 145-160.
- Goater, B., 1976. A new subspecies of *Luperina nickerlii* (Freyer) (Lep.: Noctuidae) from Cornwall. *Entomologist's Gaz.* 27: 141-144, Pl. 8.
- Hacker, H., 1989. Die Noctuidae Griechenlands Mit einer Übersicht über die Fauna des Balkansraumes (Lepidoptera: Noctuidae). *Herbipoliana*, 2. 1-589, i-xii.
- Heinicke, W. & Naumann, C., 1981. Beiträge zur Insektenfauna der DDR: Lepidoptera - Noctuidae. *Beitr. Ent.* 31(2): 395-396.
- de Joannis, J., 1925. Une nouvelle espèce de Noctuelle du genre *Palluperina* Hampson (*Luperina*, *Apamea*, auct.) appartenant au groupe de *testacea* Schiff. *Ann. Soc. Ent. Fr.* 94: 331-38; pl. 1.
- Oberthür, Ch., [1909] 1908. Observations sur une nouvelle espèce française: *Luperina graslini* [Lép. Hét.]. *Bull. Soc. ent. Fr.* 1908: 322-323.
- , 1923. Catalogue des Lépidoptères des Pyrénées-Orientales. *Et. Lép. Comp.* 20: ix-xv + 1-159pp.
- Schwingschuss, L., 1962. *Z. wien. ent. Ges.* 47: 6.
- Skinner, B., 1984. *Colour Identification Guide to Moths of the British Isles*. Viking, x + 267pp. London.
- , 1985. *Luperina nickerlii* Freyer: Sandhill Rustic in Kent. *Entomologist's Rec. J. Var.*, 97: 28.

---

#### *Coleophora deviella* Zell (Lep.: Coleophoridae) – a correction

In the short note on this species by Norman Heal (*Ent. Rec. J. Var.*, 1995, 107: 43-44) the size of the fully grown case was quoted ambitiously as being between 9 and 19mm long. Whilst such a giant *deviella* might prove attractive to entomologists, the Editor regrets that the typical case size for this species is 9 – 10mm when fully grown. A further error crept into this note. The second sentence of the second paragraph should read “. . . as I had searched this same area for several consecutive seasons I can only surmise that at the time it was a recent arrival . . .” Our apologies to Mr Heal for these errors. Paul Sokoloff, Editor.

#### An unusual foodplant for *Hedya pruniana* Hb. (Lep.: Tortricidae)

Two tortricoid larvae were obtained when beating Yew (*Taxus baccata*) at Shoreham, Kent. One spun up without feeding and emerged as, unsurprisingly, *Ditula angustiorana* Haw. The second continued to feed on *Taxus*, eventually producing an adult *Hedya pruniana*. This appears to be a considerable departure from its normal pabulum of *Prunus* and other Rosaceous trees.

– I. FERGUSON, 31 Cathcart Drive, Orpington, Kent BR6 8BU.